

**Only preparation with the  
active ingredient bibrocathol**



**Protection and disinfection  
for irritated eyes in case of  
inflamed eyelids**



# Posiformin 2%, eye ointment

**Posiformin highly suited for  
use in inflammation of the eyelids**

**The active ingredient bibrocathol contained in  
Posiformin 2% eye ointment;**

- ◆ Disinfects irritated eyelid edges and the adjacent conjunctiva
- ◆ Has an astringent (constricting) and anti-secretory effect
- ◆ Creates the conditions for quick and complication-free easing of symptoms



**Indications:**

- Unspecific, non-infected irritations of the outer eye
- Chronic inflammation of the lid margin (Blepharitis chronica)
- Non-infected fresh corneal wounds




# Posiformin 2%, eye ointment

## Successful treatment of blepharitis with bibrocathol (Posiformin® 2 %)

**Methods** In this multi-center, randomized, double-masked, placebo-controlled parallel-group comparison, the change of signs and symptoms (sum score) of blepharitis in 197 patients (ITT (intention-to-treat-group); mean age 18±56 years, 56 % female, active drug:vehicle 097:100) over 2 weeks treatment with bibrocathol 2 % eye ointment was evaluated.

**Conclusions** Blepharitis therapy with the antiseptic bibrocathol 2 % in this trial was highly efficacious and safe



### Mode of Action:

- ✓ Posiformin® 2 % contains bibrocathol which is a bismuth-containing molecule with antiseptic, astringent and secretion inhibiting properties.
- ✓ By interaction with proteins of the mucous membrane a protective membrane is formed which avoids the invasion of pathogens

Pavel A. Bezdetko & Nikolai Sergienko & Yuriy Dyomin & Andrei Korol & Nik Nikitin & Matthias Merzbacher & Dorothea Groß & Ralf Kohlen

### **Posiformin 2 %, eye ointment contains active ingredient:**

Bibrocathol The other ingredients are white soft paraffin, liquid paraffin and wool fat.

### **What Posiformin 2 % is and what it is used for?**

Pharmacotherapeutic group: Posiformin 2 % is a disinfecting, astringent and secretion inhibiting eye ointment.

### **Therapeutic indications:**

Posiformin 2 % is used in irritation of the outer eye that are not caused by bacteria, chronic (long lasting) inflammation of the eye lid margin that are not caused by bacteria and non-infectious fresh wounds of the cornea.

### **Before you use Posiformin 2 % :**

**Do not use Posiformin 2 %**

- ◆ If you are allergic to bibrocathol or any of the other ingredients of Posiformin 2 %
- ◆ While wearing contact lenses

### **Take special care with Posiformin 2 % :**

Talk to your doctor or pharmacist before using Posiformin 2 %. In persisting complaints or lack of improvement, a doctor should be consulted without delay

### **Pregnancy and breast-feeding :**

Ask your doctor or pharmacist for advice before taking any medicine.

There is not sufficient information available to establish the safety of bibrocathol in human pregnancy or lactation. Posiformin 2 % should not be used during pregnancy and lactation unless clearly necessary.

### **Dosage:**

If not otherwise prescribed by your doctor, the recommended dose for adults and children is 3 to 5 times daily a strip of about 0.5 cm of eye ointment into the conjunctival sac or on the affected position of the eye lid.

### **Possible side effects:**

Like all medicines, Posiformin 2 % can cause side effects, although not everybody gets them. These side effects include: Rare cases of allergic reactions and eye irritation with the following symptoms were reported: eye itching, eye swelling, eye pain, ocular hyperaemia, swelling of face, facial flushing. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

### **How to store Posiformin 2 % :**

Keep out of the reach and sight of children. Do not store above 25°C.

Do not use Posiformin 2 % after the expiry date which is stated on the label and carton.

The expiry date refers to the last day of that month.

**Once opened :** Posiformin 2 % should not be used longer than 4 weeks after first opening

- ✓ *Without preservative*
- ✓ *No risk of resistance development*
- ✓ *Antiseptic efficacy*
- ✓ *Excellent tolerance*

 **URSAPHARM**

**Arzneimittel GmbH**

Industriestraße 35

66129 Saarbrücken - Germany

Tel : +49(0)6805 9292-0

Fax: +49(0)6805 9292-88

info@ursapharm.de - www.ursapharm.de

